Author Index

ABADI I. *see* RODRIGUEZ M.A.
ABDUL-GAFFAR R. *see* MACINTYRE E.A.
ALBECK M. *see* SREDNI B.
AMLOT P.L. *see* BIRD P.
ANDREW E.M., ANNIS W., KAHAN M. & MAINI R.N. CD5 (Ly-1)-negative, conventional splenic B cells make a substantial contribution to the bromelain plaque-forming cell response in CBA and BW mice, 515
ANNIS W. *see* ANDREW E.M.
ARATA Y. *see* KIMURA K.

BAKIMER R. *see* MENDLOVIC S.
BALESTRA E. *see* DEL GOBBO V.
BANCHEREAU J. *see* VALLÉ A.
BARFOOT R. *see* DENHAM S.
BARRIOS H.A. *see* CAMPETELLA O.E.
BAR-SHAVIT Z. *see* LEVI-SCHAFFER F.
BARTHOLOMAEUS W.N. *see* LAWSON C.M.
BARTHOLOMEUSZ R.C.A., FORREST B.D., LABROOY J.T., EY P.L., PYLE D., SHEARMAN D.J.C. & ROWLEY D. The serum polymeric IgA antibody response to typhoid vaccination; its relationship to the intestinal IgA response, 190
BAUMANN H. *see* STADNYK A.W.
BEATTY P.G. *see* DOBRINA A.
BEDDELL C.R. *see* FRANCIS M.J.
BELL E.B. *see* WHITBY E.H.
BERANGER F. *see* SANCÉAU J.
BERTOVICH M.J. *see* NASH G.S.
BEVERLEY P.C.L. *see* WALLACE D.L.
BICKLE Q.D. *see* VIGNALI D.A.A.
BIRD P., CALVERT J.E. & AMLOT P.L. Distinctive development of IgG4 subclass antibodies in the primary and secondary responses to keyhole limpet haemocyanin in man, 355
BISHOP R. *see* GODDEERIS B.M.
BLASINI A.M. *see* RODRIGUEZ M.A.
BLEASDALE-BARR K.M. *see* DEVEY M.E.
BOLAND G.J. *see* MUDDE G.C.
BOLTON A.E. *see* POCKLEY A.G.
BOLTON E.J. *see* QUINT D.J.
BOLTON E.M. *see* MARSHALL H.E.
BONNEFOY J.-Y. *see* VALLÉ A.
BOTTAZZO G.F. *see* TODD I.
BRADLEY J. *see* NORAZMI M.N.
BRADLEY J.A. *see* MARSHALL H.E.
BRADLY B. *see* HARTNELL A.
BRENAN M. Deficient responsiveness of ovine popliteal peripheral lymphocytes, 134
BROCKE S., DAYAN M., ROTHBARD J., FUCHS S. & MOZES E. The autoimmune response of different mouse strains to T-cell epitopes of the human acetylcholine receptor α subunit, 495
BROCKE S. *see* MENDLOVIC S.
BROWN A.L. *see* FRANCIS M.J.
BROWN F. *see* FRANCIS M.J.
BRUIJNZEEL P.L.B. *see* MUDDE G.C.
BRUIJNZEEL-KOOMEN C.A.F.M. *see* MUDDE G.C.
BUCKHAM S.P. *see* QUINT D.J.
BUJDOSO R., YOUNG P., HOPKINS J. & MCCONNELL I. IL-2-like activity in lymph fluid following *in vivo* antigen challenge, 45
BURKART V. *see* ZIELASEK J.
BUTCHER R.D.J., MCGARVIE G.M. & CUSHLEY W. Recombinant interleukin-4 promotes expression of the CD25 (Tac) antigen at the plasma membrane of high-density human tonsillar B lymphocytes, 57

CALIO R. *see* DEL GOBBO V.
CALVERT J.E. *see* BIRD P.
CAMPETELLA O.E., GALASSI N.V. & BARRIOS H.A. Contrasuppressor cells induced by Junin virus, 629
CANTRELL D. *see* FOXWELL B.M.J.
CAPRON A. *see* HAQUE A.
CARLOS T.M. *see* DOBRINA A.
CARRIER M. *see* LIEW F.Y.
CARTER D.B. *see* SANCÉAU J.
CHAMEKH M. *see* HAQUE A.
CHAMPION B.R. *see* QUINT D.J.
CHANG H.R., GRAU G.E. & PECHÈRE J.C. Role of TNF and IL-1 in infections with *Toxoplasma gondii*, 33
CHATURVEDI U.C. *see* KHANNA M.
CHILDERSTONE A., HARON J. & LEHNER T. The reactivity of naturally sensitized human CD4 cells and IgG antibodies to synthetic peptides derived from the amino terminal sequences of a 3800 MW *Streptococcus mutans* antigen, 177
CHOY W.F. *see* LIM P.L.
CHRISTIE J.F., DUNBAR B., DAVIDSON I. & KENNEDY M.W. N-terminal amino acid sequence identity between a major allergen of *Ascaris lumbricoides* and *Ascaris suum*, and MHC-restricted IgE responses to it, 596
CHUA T.P., SMITH C.E., REITH R.W. & WILLIAMSON J.D. Inflammatory responses and the generation of chemoattractant activity in cowpox virus-infected tissues, 202
CLARKE B.E. *see* FRANCIS M.J.
CLAYTON J., RHIND S.M., GROVES D.J. & MORRIS B.A. Pharmacokinetic studies of passively administered ovine anti-testosterone antibodies given to cattle by the subcutaneous and intravenous routes, 603
COCKER J.E. *see* MARSHALL H.E.
COLEMAN J.W. Conditioned medium from concanavalin A-stimulated spleen cells inhibits the IgE-dependent sensitization of murine peritoneal mast cells *in vitro*, 150
ČOLIĆ M., POPOVIĆ L.J., GAŠIĆ S., DRAGOJEVIĆ-SIMIĆ V., MILIĆEVIĆ N.M., MATANOVIĆ D. & DUJIĆ A. Immunohistochemical characterization of rat thymic non-lymphoid cells. II. Macrophages and granulocytes defined by monoclonal antibodies, 416
COLLEN T. *see* TAKAMATSU H.
COLOTTA F. *see* SICA A.
CORNELIS J. *see* HAQUE A.
CRAMER R. *see* ZABUCCHI G.
CROMWELL O. *see* MOQBEL R.
CUSHLEY W. *see* BUTCHER R.D.J.
CZUPRYNSKI C.J. *see* ROLL J.T.

DAL CANTO M.C. *see* DEVEY M.E.
DANIELS R.H., HOUSTON W.A.J., PETERSEN M.M., WILLIAMS J.D., WILLIAMS B.D. & MORGAN B.P. Stimulation of human rheumatoid synovial cells by non-lethal complement membrane attack, 237
DAVIDSON I. *see* CHRISTIE J.F.
DAWSON R. *see* MIYATA T.
DAYAN M. *see* BROCKE S.
DE ANDRES B., DEL POZO V., MARTIN E., PALOMINO P. & LAHOZ C. Release of O_2^- and LTC_4 by murine eosinophils: role of intra- and extracellular calcium, 271
DE GAST G.C. *see* MUDDE G.C.
DE PANFILIS G., MANARA G.C., FERRARI C., TORRESANI C. & ROWDEN G. Subsets of keratinocytes and Langerhans' cells express epitopes associated with suppressor-inducer capabilities in resting normal human epidermis, 622

DEJANA E. *see* SICA A.
DEL GOBBO V., VILLANI N., MARINI S., BALESTRA E. & CALIO R. Suppressor cells induced by influenza virus inhibit interleukin-2 production in mice, 454
DEL POZO V. *see* DE ANDRES B.
DE SANCTIS J.B. *see* RODRIGUEZ M.A.
DENHAM S., BARFOOT R. & SILLS J. The rat RcR for monomeric IgG is preferentially expressed on red pulp macrophages in the spleen, 329
DENYER M.S. *see* TAKAMATSU H.
DEVEY M.E., MAJOR P.J., BLEASDALE-BARR K.M., HOLLAND G.P., DAL CANTO M.C. & PATERSON P.Y. Experimental allergic encephalomyelitis (EAE) in mice selectively bred to produce high affinity (HA) or low affinity (LA) antibody responses, 519
DEVEY M.E. *see* VIGNALI D.A.A.
DIAZ-SANCHEZ D. & KEMENY D.M. The sensitivity of rat $CD8^+$ and $CD4^+$ T cells to ricin *in vivo* and *in vitro* and their relationship to IgE regulation, 71
DIMARCHI R. *see* FLYNN J.N.
DOBRINA A., CARLOS T.M., SCHWARTZ B.R., BEATTY P.G., OCHS H.D. & HARLAN J.M. Phorbol ester causes down-regulation of CD11/CD18-independent neutrophil adherence to endothelium, 429
DOEL T. *see* FLYNN J.N.
DORRINGTON K.J. *see* WILLIAMS D.B.
DOUGAN G. *see* O'CALLAGHAN D.
DRAGOJEVIĆ-SIMIĆ V. *see* ČOLIĆ M.
DUJIĆ A. *see* ČOLIĆ M.
DUNBAR B. *see* CHRISTIE J.F.

EY P.L. *see* BARTHOLOMEUSZ R.C.A.

FAGER G. *see* STEMME S.
FALCOFF R. *see* SANCÉAU J.
FELDMANN M. *see* TODD I.
FERRARI C. *see* DE PANFILIS G.
FERREIRA O.C. *see* GARCIA-PARDO A.
FERRO L.M. & ZOLA H. Modulation of expression of the antigen identified by FMC7 upon human B-lymphocyte activation: evidence for differences between activation *in vivo* and *in vitro*, 373
FLEISCHER B., FLEISCHER S., MAIER K., WIEDMANN K.H., SACHER M., THALER H. & VALLBRACHT A. Clonal analysis of infiltrating T lymphocytes in liver tissue in viral hepatitis A, 14
FLEISCHER S. *see* FLEISCHER B.
FLORES-ROMO L., MILLSUM M.J., GILLIS S., STUBBS P., SYKES C. & GORDON J. Immunoglobulin isotype production by cycling human B lymphocytes in response to recombinant cytokines and anti-IgM, 342
FLYNN J.N., HARKISS G.D., DOEL T. & DIMARCHI R. Analysis of immune responses in the sheep to synthetic peptides of foot-and-mouth disease virus using ovine polyclonal and monoclonal antibodies, 1
FORREST B.D. *see* BARTHOLOMEUSZ R.C.A.
FOXWELL B.M.J., SIMON J., HERRERO J.-J., TAYLOR D., WOERLY G., CANTRELL D. & RYFFEL B. Anti-CD3 antibody-induced expression of both p55 and p75 chains of the high-affinity interleukin-2 receptor on human T lymphocytes is inhibited by cyclosporin A, 104
FRANCIS M.J., HASTINGS G.Z., CLARKE B.E., BROWN A.L., BEDDELL C.R., ROWLANDS D.J. & BROWN F. Neutralizing antibodies to all seven serotypes of foot-and-mouth disease virus elicited by synthetic peptides, 171
FREEDMAN A.S. *see* VALLÉ A.
FREEMAN G. *see* VALLÉ A.
FRICKE H. *see* MENDLOVIC S.
FUCHS S. *see* BROCKE S.
FUJIWARA M. *see* MATSUMOTO Y.

GALASSI N.V. *see* CAMPETELLA O.E.
GARCIA-PARDO A. & FERREIRA O.C. Adhesion of human T-lymphoid cells to fibronectin is mediated by two different fibronectin domains, 121
GARRONE P. *see* VALLÉ A.
GAŠIĆ S. *see* ČOLIĆ M.
GAULDIE J. *see* STADNYK A.W.
GILLIS S. *see* FLORES-ROMO L.
GODDEERIS B.M., MORRISON W.I., TOYE P.G. & BISHOP R. Strain specificity of bovine *Theileria parva*-specific cytotoxic T cells is determined by the phenotype of the restricting class I MHC, 38
GOLDSCHMIDT T.J., JANSSON L. & HOLMDAHL R. *In vivo* elimination of T cells expressing specific T-cell receptor $V\beta$ chains in mice susceptible to collagen-induced arthritis, 508
GOLDSTEIN A. *see* ZIELASEK J.
GORDON J. *see* FLORES-ROMO L.
GRACIE J.A. *see* MARSHALL H.E.
GRAU G.E. *see* CHANG H.R.
GREENWOOD J.H. *see* WOOD G.W.
GROVES D.J. *see* CLAYTON J.

HALLIDAY G.M., WOOD R.C. & MULLER H.K. Presentation of antigen to suppressor cells by a dimethylbenz (*a*) anthracene-resistant, Ia-positive, Thy-1-negative, I-J-restricted epidermal cell, 97
HAMADA S. *see* OGAWA T.
HAMMOND L.J. *see* TODD I.
HANSSON G.K. *see* STEMME S.
HAQUE A., CHAMEKH M., CORNELIS J., CAPRON A. & HAQUE S. A monoclonal antibody to Ly-6 gene product inhibits generation of functionally active T cells and recognizes single antigenic specificity whose expression is up-regulated in virus-transformed rat fibroblast, 558
HAQUE S. *see* HAQUE A.
HARKISS G.D. *see* FLYNN J.N.
HARLAN J.M. *see* DOBRINA A.
HARON J. *see* CHILDERSTONE A.
HART D.N.J. *see* PRICKETT T.C.R.
HARTNELL A., MOQBEL R., WALSH G.M., BRADLY B. & KAY A.B. Fcγ and CD11/CD18 receptor expression on normal density and low density human eosinophils, 264
HARVEY J., JONES D.B. & WRIGHT D.H. Differential expression of MHC- and macrophage-associated antigens in human fetal and postnatal small intestine, 409
HASEGAWA M. *see* MIYATA T.
HASKARD D.O. *see* THORNHILL M.H.
HASTINGS G.Z. *see* FRANCIS M.J.
HERRERO J.-J. *see* FOXWELL B.M.J.
HIDESHIMA T., OKADA N. & OKADA H. Expression of HRF20, a regulatory molecule of complement activation, on peripheral blood mononuclear cells, 396
HINES D.L. *see* JOHNSON L.L.
HOHMANN A.W. *see* NORAZMI M.N.
HOLÁŇ V. & LIPOLDOVÁ M. Tumour-bearing animals synthesize a decreased level of mRNA for the inducible 55,000 MW interleukin-2 receptor, 626
HOLLAND G.P. *see* DEVEY M.E.
HOLMDAHL R. *see* GOLDSCHMIDT T.J.
HOLMES M.J. *see* RAMSAY A.J.
HOLT P.G. *see* NELSON D.
HOPKINS J. *see* BUJDOSO R.
HOSTETTER M.K. *see* MOY J.N.
HOUSTON W.A.J. *see* DANIELS R.H.
HOWARD C.J. *see* NAESSENS J.
HUNT S.V. *see* VONDERHEIDE R.H.
HUNTLEY J.F. *see* IRVINE J.

IKEZAWA H. *see* MIYATA T.
INAGI R. *see* MIYATA T.
INOUE I. *see* MIYATA T.
IRVINE J., NEWLANDS G.F.J., HUNTLEY J.F. & MILLER H.R.P. Interaction of murine intestinal mast cell proteinase with inhibitors (serpins) in blood; analysis by SDS–PAGE and Western blotting, 139
ISENBERG D.A. *see* WATTS R.A.
ISOBE K. *see* MIYATA T.
ISSEKUTZ A.C. *see* MEGYERI P.
ISSEKUTZ T.B. *see* MEGYERI P.
ITO M. *see* YAMAMOTO K.

JAMES R.F.L. *see* TODD I.
JANSSON L. *see* GOLDSCHMIDT T.J.
JARVIS L.R. *see* NORAZMI M.N.
JOHNSON L.L. & HINES D.L. Tumour-induced immunity to H-Y-disparate skin grafts without concomitant priming of CTL, 85

JONES B. *see* NAESSENS J.
JONES D.B. *see* HARVEY J.
JONES D.B. *see* NESBITT A.M.

KAHAN M. *see* ANDREW E.M.
KALECHMAN Y. *see* SREDNI B.
KAWAI K. *see* MATSUMOTO Y.
KAY A.B. *see* HARTNELL A.
KAY A.B. *see* MOQBEL R.
KEMENY D.M. *see* DIAZ-SANCHEZ D.
KENNEDY M.W. *see* CHRISTIE J.F.
KHANNA M., CHATURVEDI U.C., SHARMA M.C., PANDEY V.C. & MATHUR A. Increased capillary permeability mediated by a dengue virus-induced lymphokine, 449
KIESEL U. *see* ZIELASEK J.
KIMOTO M. *see* MINETA T.
KIMURA K., ARATA Y., YASUDA T., KINOSITA K. & NAKANISHI M. Location of membrane-bound hapten with different length spacers, 323
KINOSITA K. *see* KIMURA K.
KLEIN M.H. *see* WILLIAMS D.B.
KOBAYASHI S. *see* MIYATA T.
KOLB H. *see* ZIELASEK J.
KOUASSI E. *see* LI Y.S.
KULCZYCKI A. *see* NASH G.S.
KUROSAWA Y. *see* YAMAUCHI T.
KURTZ R.S. *see* ROLL J.T.
KUSUMOTO Y. *see* OGAWA T.
KYAN-AUNG U. *see* THORNHILL M.H.

LABROOY J.T. *see* BARTHOLOMEUSZ R.C.A.
LAHOZ C. *see* DE ANDRES B.
LAWSON C.M., O'DONOGHUE H., BARTHOLOMAEUS W.N. & REED W.D. Genetic control of mouse cytomegalovirus-induced myocarditis, 20
LECK P.A. *see* MOWAT A. MCI.
LEE T.H. *see* THORNHILL M.H.
LEHNER T. *see* CHILDERSTONE A.
LEON-PONTE M. *see* RODRIGUEZ M.A.
LEVI-SCHAFFER F. & BAR-SHAVIT Z. Osteoblast-like cell line maintains *in vitro* rat peritoneal mast cell viability and functional activity, 145
LEWIS C.E., MCCARTHY S.P., LORENZEN J. & MCGEE J.O'D. Differential effects of LPS, IFN-γ and TNFα on the secretion of lysozyme by individual human mononuclear phagocytes: relationship to cell maturity, 402
LI Y.S., KOUASSI E. & REVILLARD J.P. Differential regulation of mouse B-cell activation by β-adrenoceptor stimulation depending on type of mitogens, 367
LIEW F.Y., PARKINSON C., MILLOTT S., SEVERN A. & CARRIER M. Tumour necrosis factor (TNFα) in leishmaniasis. I. TNFα mediates host protection against cutaneous leishmaniasis, 570
LIM P.L. & CHOY W.F. A thymus-dependent (type 1) phosphorylcholine antigen isolated from *Trichinella spiralis* protects mice against pneumococcal infection, 443
LINCH D.C. *see* MACINTYRE E.A.
LIPOLDOVÁ M. *see* HOLÁŇ V.
LORENZEN J. *see* LEWIS C.E.

MCCARTHY S.P. *see* LEWIS C.E.
MCCONNELL I. *see* BUJDOSO R.
MACDERMOTT R.P. *see* NASH G.S.
MACDONALD A.J. *see* MOQBEL R.
MCGARVIE G.M. *see* BUTCHER R.D.J.
MCGEE J.O'D. *see* LEWIS C.E.
MCHUGH N. *see* NAESSENS J.
MACINTYRE E.A., ROBERTS P.J., ABDUL-GAFFAR R., MORGAN J. & LINCH D.C. Activation of monocytic cells by monoclonal antibodies to the CD11a/18 (LFA-1) complex: mediation by Fc receptor, 574
MAEDA K. *see* MIYATA T.
MAIER K. *see* FLEISCHER B.
MAINI R.N. *see* ANDREW E.M.
MAJOR P.J. *see* DEVEY M.E.
MANARA G.C. *see* DE PANFILIS G.
MANTOVANI A. *see* SICA A.
MARINI S. *see* DEL GOBBO V.
MARSHALL H.E., BOLTON E.M., GRACIE J.A., COCKER J.E., SANDILANDS G.P. & BRADLEY J.A. FcR blocking activity in serum of actively enhanced rat renal allograft recipients is due to IgG anti-class II MHC alloantibody, 379
MARTIN E. *see* DE ANDRES B.
MASKELL D. *see* O'CALLAGHAN D.
MATANOVIĆ D. *see* ČOLIĆ M.
MATHUR A. *see* KHANNA M.
MATSUMOTO K. *see* OGIMOTO M.
MATSUMOTO Y., KAWAI K., TOMITA Y. & FUJIWARA M. Limiting-dilution analysis of the frequency of myelin basic protein-reactive T cells in Lewis, PVG/c and BN rats. Implication for susceptibility to autoimmune encephalomyelitis, 215
MATSUNAGA M. *see* MINETA T.
MATSUSHIMA K. *see* SICA A.
MATSUZAKI G. *see* OGIMOTO M.
MAUSER L. *see* WOOD G.W.
MEGYERI P., SADOWSKA J., ISSEKUTZ T.B. & ISSEKUTZ A.C. Endotoxin-stimulated human macrophages produce a factor that induces polymorphonuclear leucocyte infiltration and is distinct from interleukin-1, tumour necrosis factor α and chemotactic factors, 155
MENDLOVIC S., BROCKE S., FRICKE H., SHOENFELD Y., BAKIMER R. & MOZES E. The genetic regulation of the induction of experimental SLE, 228
MENEGAZZI R. *see* ZABUCCHI G.
MILIĆEVIĆ N.M. *see* ČOLIĆ M.
MILLER H.R.P. *see* IRVINE J.
MILLOTT S. *see* LIEW F.Y.
MILLSUM M.J. *see* FLORES-ROMO L.
MINETA T., SEKI K., MATSUNAGA M. & KIMOTO M. Existence of mixed isotype $A_\beta E_\alpha$ class II molecules in E^d_α gene-introduced C57BL/6 transgenic mice, 385
MIYAMOTO T. *see* YAMAMOTO K.
MIYATA T., ISOBE K., DAWSON R., RITTER M.A., INAGI R., ODA O., TAGUCHI R., IKEZAWA H., INOUE I., SEO H., HASEGAWA M., KOBAYASHI S., MAEDA K., YAMADA K. & NAKASHIMA I. Determination of the molecular nature and cellular localization of Thy-1 in human renal tissue, 391
MOORE K. *see* NESBITT A.M.
MOQBEL R., MACDONALD A.J., CROMWELL O. & KAY A.B. Release of leukotriene C_4 (LTC_4) from human eosinophils following adherence to IgE- and IgG-coated schistosomula of *Schistosoma mansoni*, 435
MOQBEL R. *see* HARTNELL A.
MORGAN B.P. *see* DANIELS R.H.
MORGAN J. *see* MACINTYRE E.A.
MORI A. *see* YAMAMOTO K.
MORRIS B.A. *see* CLAYTON J.
MORRISON W.I. *see* GODDEERIS B.M.
MOWAT A. MCI. & LECK P.A. Generalized toxicity of L-leucyl-leucine-methyl ester for lymphocyte functions, 564
MOY J.N., NELSON R.D., RICHARDS K.L. & HOSTETTER M.K. Identification of an IgA inhibitor of neutrophil chemotaxis and its membrane target for the metabolic burst, 257
MOZES E. *see* BROCKE S.
MOZES E. *see* MENDLOVIC S.
MUDDE G.C., VAN REIJSEN F.C., BOLAND G.J., DE GAST G.C., BRUIJNZEEL P.L.B. & BRUIJNZEEL-KOOMEN C.A.F.M. Allergen presentation by epidermal Langerhans' cells from patients with atopic dermatitis is mediated by IgE, 335
MULLER H.K. *see* HALLIDAY G.M.
MURPHY J.J., TRACZ M. & NORTON J.D. Patterns of nuclear proto-oncogene expression during induced differentiation and proliferation of human B chronic lymphocytic leukaemia cells, 490

NADLER L.M. *see* VALLÉ A.
NAESSENS J., NEWSON J., MCHUGH N., HOWARD C.J., PARSONS K. & JONES B. Characterization of a bovine leucocyte differentiation antigen of 145,000 MW restricted to B lymphocytes, 525
NAKAHAMA T. *see* YAMAMOTO K.
NAKANISHI M. *see* KIMURA K.
NAKANO K. *see* OKAMOTO H.
NAKASHIMA I. *see* MIYATA T.
NAOE T. *see* YAMAUCHI T.
NARDON E. *see* ZABUCCHI G.

NASH G.S., MACDERMOTT R.P., SCHLOEMANN S., BERTOVICH M.J., O'NEAL J., PORTER L. & KULCZYCKI A. Bovine IgG1, but not IgG2, binds to human B cells and inhibits antibody secretion, 361
NAYLOR P. *see* ZIELASEK J.
NELSON D., STRICKLAND D. & HOLT P.G. Selective attrition of non-recirculating T cells during normal passage through the lung vuscular bed, 476
NELSON R.D. *see* MOY J.N.
NESBITT A.M., JONES D.B. & MOORE K. Phenotypic changes in a CD4+ lymphocyte subset correlate with a conversion from a suppressor to helper inducer function, 65
NEWLANDS G.F.J. *see* IRVINE J.
NEWSON J. *see* NAESSENS J.
NOMOTO K. *see* OGIMOTO M.
NORAZMI M.N., HOHMANN A.W., SKINNER J.M., JARVIS L.R. & BRADLEY J. Density and phenotype of tumour-associated mononuclear cells in colonic carcinomas determined by computer-assisted video image analysis, 282
NORTON J.D. *see* MURPHY J.J.

O'CALLAGHAN D., MASKELL D., TITE J. & DOUGAN G. Immune responses in BALB/c mice following immunization with aromatic compound or purine-dependent *Salmonella typhimurium* strains, 184
O'DONOGHUE H. *see* LAWSON C.M.
O'NEAL J. *see* NASH G.S.
OCHS H.D. *see* DOBRINA A.
ODA O. *see* MIYATA T.
OGAWA T., SHIMAUCHI H., KUSUMOTO Y. & HAMADA S. Humoral immune response to *Bacteroides gingivalis* fimbrial antigen in mice, 8
OGIMOTO M., YOSHIKAI Y., MATSUZAKI G., OHGA S., MATSUMOTO K. & NOMOTO K. Clonal deletion of self-Mls-reactive thymocyts at the early stage in H-2-compatible but Mls-disparate radiation chimeras, 482
OHGA S. *see* OGIMOTO M.
OHNO I., TANNO Y., YAMAUCHI K. & TAKISHIMA T. Production of tumour necrosis factor by mastocytoma P815 cells, 312
OKADA H. *see* HIDESHIMA T.
OKAKA N. *see* HIDESHIMA T.
OKAMOTO H. & NAKANO K. Regulation of interleukin-1 synthesis by histamine produced by mouse peritoneal macrophages *per se*, 162
OKUDAIRA H. *see* YAMAMOTO K.
OLDSTONE M.B.A. *see* SHYP S.

PALOMINO P. *see* DE ANDRES B.
PANDEY V.C. *see* KHANNA M.
PARKHOUSE R.M.E. Three B-cell surface molecules associating with membrane immunoglobulin, 298
PARKINSON C. *see* LIEW F.Y.
PARSONS K. *see* NAESSENS J.
PATERSON P.Y. *see* DEVEY M.E.
PATRIARCA P. *see* ZABUCCHI G.
PAWELEC G. CD4+ CD45R+ suppressor-inducer T-cell clones: requirements for cellular interaction, proliferation and lymphokines for the induction of suppression in peripheral blood mononuclear cells, 536
PECHÈRE J.C. *see* CHANG H.R.
PERERA M.A. *see* WILLIAMS D.B.
PETERSEN M.M. *see* DANIELS R.H.
POCKLEY A.G. & BOLTON A.E. The effect of human placental protein 14 (PP14) on the production of interleukin-1 from mitogenically stimulated mononuclear cell cultures, 277
POLENTARUTTI N. *see* SICA A.
POPOVIĆ L.J. *see* ČOLIĆ M.
PORTER L. *see* NASH G.S.
PRICKETT T.C.R. & HART D.N.J. Anti-leucocyte common (CD45) antibodies inhibit dendritic cell stimulation of CD4 and CD8 T-lymphocyte proliferation, 250
PYLE D. *see* BARTHOLOMEUSZ R.C.A.

QUINT D.J., BUCKHAM S.P., BOLTON E.J., SOLARI R., CHAMPION B.R. & ZANDERS E.D. Immunoregulation in the common marmoset, *Calithrix jaccus*: functional properties of T and B lymphocytes and their response to human interleukins 2 and 4, 616

RAMSAY A.J. & HOLMES M.J. A rabbit model for mucosal immunity in the bowel. II. Local cellular reactivity to virus infection, 27
RAVIRAJAN C.T. *see* WATTS R.A.
REED W.D. *see* LAWSON C.M.
REITH R.W. *see* CHUA T.P.
REVILLARD J.P. *see* LI Y.S.
RHIND S.M. *see* CLAYTON J.
RICHARDS K.L. *see* MOY J.N.
RIFAI A. & WU C.-G. Clearance kinetics, tissue localization and fate of IgA–anti-idiotype complexes, 610
RITTER M.A. *see* MIYATA T.
ROBERTS P.J. *see* MACINTYRE E.A.
RODRIGUEZ M.A., DE SANCTIS J.B., BLASINI A.M., LEON-PONTE M. & ABADI I. Human IFN-γ up-regulates IL-2 receptors in mitogen-activated T lymphocytes, 554
ROLL J.T., YOUNG K.M., KURTZ R.S. & CZUPRYNSKI C.J. Human rTNFα augments anti-bacterial resistance in mice: potentiation of its effects by recombinant human rIL-1α, 316
ROTHBARD J. *see* BROCKE S.
ROWDEN G. *see* DE PANFILIS G.
ROWLANDS D.J. *see* FRANCIS M.J.
ROWLEY D. *see* BARTHOLOMEUSZ R.C.A.
RYFFEL B. *see* FOXWELL B.M.J.

SACHER M. *see* FLEISCHER B.
SADOWSKA J. *see* MEGYERI P.
SANCÉAU J., FALCOFF R., BERANGER F., CARTER D.B. & WIETZERBIN J. Secretion of interleukin-6 (IL-6) by human monocytes stimulated by muramyl dipeptide and tumour necrosis factor alpha, 52
SANDILANDS G.P. *see* MARSHALL H.E.
SCHICK B. Cleavage of a rat serosal mast cell membrane component during degranulation mediated by chymase, a secretory granule protease, 423
SCHLOEMANN S. *see* NASH G.S.
SCHMIDT R.E. *see* WITTE T.
SCHWARTZ B.R. *see* DOBRINA A.
SEKI K. *see* MINETA T.
SEO H. *see* MIYATA T.
SEVERN A. *see* LIEW F.Y.
SHALIT F. *see* SREDNI B.
SHARMA M.C. *see* KHANNA M.
SHEARMAN D.J.C. *see* BARTHOLOMEUSZ R.C.A.
SHIKU H. *see* YAMAUCHI T.
SHIMAUCHI H. *see* UGAWA T.
SHOENFELD Y. *see* MENDLOVIC S.
SHYP S., TISHON A. & OLDSTONE M.B.A. Inhibition of diabetes in BB rats by virus infection. II. Effect of virus infection on the immune response to non-viral and viral antigens, 501
SICA A., MATSUSHIMA K., VAN DAMME J., WANG J.M., POLENTARUTTI N., DEJANA E., COLOTTA F. & MANTOVANI A. IL-1 transcriptionally activates the neutrophil chemotactic factor/IL-8 gene in endothelial cells, 548
SILLS J. *see* DENHAM S.
SIMON J. *see* FOXWELL B.M.J.
SKINNER J.M. *see* NORAZMI M.N.
SMITH C.E. *see* CHUA T.P.
SOLARI R. *see* QUINT D.J.
SPARSHOTT S.M. *see* WHITBY E.H.
SREDNI B., KALECHMAN Y., SHALIT F. & ALBECK M. Synergism between AS101 and PMA in lymphokine production, 110
STADNYK A.W., BAUMANN H. & GAULDIE J. The acute-phase protein response in parasite infection. *Nippostrongylus brasiliensis* and *Trichinella spiralis* in the rat, 588
STAINES N.A. *see* WATTS R.A.
STEMME S., FAGER G. & HANSSON G.K. MHC class II antigen expression in human vascular smooth muscle cells is induced by interferon-gamma and modulated by tumour necrosis factor and lymphotoxin, 243
STRICKLAND D. *see* NELSON D.
STUBBS P. *see* FLORES-ROMO L.
SYKES C. *see* FLORES-ROMO L.

TAGUCHI R. *see* MIYATA T.
TAKAMATSU H., COLLEN T., TUCKER D.L. & DENYER M.S. Characteriza-

tion of long-term cultured bovine CD4-positive and CD8-positive T-cell lines and clones, 293
TAKISHIMA T. *see* OHNO I.
TANNO Y. *see* OHNO I.
TAYLOR D. *see* FOXWELL B.M.J.
TAYLOR M.G. *see* VIGNALI D.A.A.
THALER H. *see* FLEISCHER B.
THORNHILL M.H., KYAN-AUNG U., LEE T.H. & HASKARD D.O. The cells and neutrophils exhibit differential adhesion to cytokine-stimulated endothelial cells, 287
TISHON A. *see* SHYP S.
TITE J. *see* O'CALLAGHAN D.
TODD I., HAMMOND L.J., JAMES R.F.L., FELDMANN M. & BOTTAZZO G.F. Epidermal growth factor and transforming growth factor-alpha suppress HLA class II induction in human thyroid epithelial cells, 91
TOMITA Y. *see* MATSUMOTO Y.
TORRESANI C. *see* DE PANFILIS G.
TOYE P.G. *see* GODDEERIS B.M.
TRACZ M. *see* MURPHY J.J.
TUCKER D.L. *see* TAKAMATSU H.

VALLBRACHT A. *see* FLEISCHER B.
VALLÉ A., GARRONE P., YSSEL H., BONNEFOY J.-Y., FREEDMAN A.S., FREEMAN G., NADLER L.M. & BANCHEREAU J. mAb 104, a new monoclonal antibody, recognizes the B7 antigen that is expressed on activated B cells and HTLV-1-transformed T cells, 531
VAN DAMME J. *see* SICA A.
VAN REIJSEN F.C. *see* MUDDE G.C.
VIGNALI D.A.A., DEVEY M.E., BICKLE Q.D. & TAYLOR M.G. The role of antibody affinity and titre in immunity to *Schistosoma mansoni* following vaccination with highly irradiated cercariae, 195
VILLANI N. *see* DEL GOBBO V.
VONDERHEIDE R.H. & HUNT S.V. Does the availability of either B cells or $CD4^+$ cells limit germinal centre formation? 487
VONDERHEIDE R.H. & HUNT S.V. T lymphocytes in rat germinal centres belong to an $ER3^+$ subpopulation of $CD4^+$ cells, 542

WALLACE D.L. & BEVERLEY P.C.L. Phenotypic changes associated with activation of $CD45RA^+$ and $CD45RO^+$ T cells, 460
WANG J.M. *see* SICA A.
WATTS R.A., RAVIRAJAN C.T., STAINES N.A. & ISENBERG D.A. A human fetal monoclonal DNA-binding antibody shares idiotypes with fetal and adult murine monoclonal DNA-binding antibodies, 348
WHITBY E.H., SPARSHOTT S.M. & BELL E.B. Allograft rejection in athymic nude rats by transferred T-cell subsets. I. The response of naive $CD4^+$ and $CD8^+$ thoracic duct lymphocytes to complete allogeneic incompatabilities, 78
WIEDMANN K.H. *see* FLEISCHER B.
WIETZERBIN J. *see* SANCÉAU J.
WILKINSON P.C. Relation between locomotion, chemotaxis and clustering of immune cells, 127
WILLIAMS B.D. *see* DANIELS R.H.
WILLIAMS D.B., PERERA M.A., DORRINGTON K.J. & KLEIN M.H. Respective contribution of intracellular calcium release and extracellular calcium influx for interleukin-2 synthesis in activated T-cell hybrids, 468
WILLIAMS J.D. *see* DANIELS R.H.
WILLIAMSON J.D. *see* CHUA T.P.
WITTE T., WORDELMANN K. & SCHMIDT R.E. Heterogeneity of human natural killer cells in the spleen, 166
WALSH G.M. *see* HARTNELL A.
WOERLY G. *see* FOXWELL B.M.J.
WOOD G.W., GREENWOOD J.H. & MAUSER L. Growth of murine thymocytes *in vitro* in chemically defined medium, 303
WOOD R.C. *see* HALLIDAY G.M.
WORDELMANN K. *see* WITTE T.
WRIGHT D.H. *see* HARVEY J.
WU C.-G. *see* RIFAI A.

YAMADA K. *see* MIYATA T.
YAMADA K. *see* YAMAUCHI T.
YAMAMOTO K., MORI A., NAKAHAMA T., ITO M., OKUDAIRA H. & MIYAMOTO T. Experimental treatment of autoimmune MRL-lpr/lpr mice with immunosuppressive compound FK506, 222
YAMAUCHI K. *see* OHNO I.
YAMAUCHI T., NAOE T., KUROSAWA Y., SHIKU H. & YAMADA K. Autoantibodies to c-*myc* nuclear protein products in autoimmune disease, 117
YASUDA T. *see* KIMURA K.
YOSHIKAI Y. *see* OGIMOTO M.
YOUNG K.M. *see* ROLL J.T.
YOUNG P. *see* BUJDOSO R.
YSSEL H. *see* VALLÉ A.

ZABUCCHI G., MENEGAZZI R., CRAMER R., NARDON E. & PATRIARCA P. Mutal influence between eosinophil preoxidase (EPO) and neutrophils: neutrophils reversibly inhibit EPO enzymatic and EPO increases neutrophil adhesiveness, 580
ZANDERS E.D. *see* QUINT D.J.
ZIELASEK J., BURKART V., NAYLOR P., GOLDSTEIN A., KIESEL U. & KOLB H. Interleukin-2-dependent control of disease development in spontaneously diabetic BB rats, 209
ZOLA H. *see* FERRO L.M.

Subject Index

Acute-phase protein response:
in nematode infection, 588
Adhesion of human T cells to fibronectin:
mediated by two different fibronectin domains, 121
Adhesion of T cells and polymorphs to endothelial cells, 287
Affinity and titre of antibody:
role of *S. mansoni* immunity, 195
Allograft rejection:
by T-cell subsets, 78
Antigen presentation, by epidermal Langerhans' cells:
in atopic dermatitis, 335
Antigen presentation, to suppressor cells, by epidermal cells, 97
Anti-idiotype complexes:
IgA, clearance kinetics, localization and fate, 610
Anti-testosterone antibodies:
pharmacokinetics, 603
Autoantibodies:
to c-*myc* products, 117
Autoimmune response:
to T-cell epitopes of the acetylcholine receptor, 495

B7 antigen:
on activated B cells and HTLV-1-transformed T cells, 531
B-cell activation:
β-adrenergic regulation, 367
B-cell surface molecules:
associating with membrane Ig, 298
B-lymphocyte activation:
modulation of FMC7 antigen, 373
B lymphocytes:
oncogene expression, 490
effect of cytokines and anti-IgM on isotype production, 342
Bovine B-cell antigen, 525
Bovine $CD4^+$ and $CD8^+$ T-cell lines and clones, 293
Bovine immunoglobulin:
effects of human B cells, 361

CD4 cells:
natural reactivity of *Streptococcus mutans* synthetic peptides, 177
$CD4^+$ lymphocyte subset:
phenotypic changes correlate with conversion from suppressor to helper-inducer function, 65
$CD45RA^+$ and $CD45RO^+$ T cells:
phenotypic changes associated with activation, 460
CD5-negative B cells:
role in bromelain PFC response, 515
Chemoattractant activity:
and cowpox viruses, 202
Chemotaxis, neutrophils:
inhibition by IgA, 257
Class II induction:
suppression by epidermal growth factor and transforming factor-alpha, 91
Clonal deletion of self-Mls-reactive thymocytes:
in radiation chimeras, 482
Cyclosporin A:
inhibition of anti-CD3-induced IL-2 receptor expression, 104
Cytomegalovirus:
induction of myocarditis, genetic control, 20
Cytotoxic T cells:
T. parva-specific, determination of specificity by MHC class I, 38

Dendritic cell stimulation of CD4 and CD8:
inhibition by anti-CD45, 250
Dengue virus:
increased capillary permeability induced by lymphokines, 449
Diabetes:
inhibition by virus infection, 501
Diabetes, autoimmune:
control by IL-2, 209
DY-induced suppression, 536

Encephalomyelitis, autoimmune:
role of MBP-reactive T cells, 215
Eosinophil peroxidase:
neutrophil interaction, 580
Eosinophils:
Fcγ and CD11/CD18 receptor expression, 264
release of O_2^- and LTC_4, role of calcium, 271
Epidermal cell:
antigen presentation to suppressor cells, 97
Epidermal growth factor, and transforming growth factor-alpha:
suppression of class II, 91
Exotoxin-stimulated human macrophages:
production of factor inducing PMN infiltration, 155
Experimental allergic encephalomyelitis:
in high-affinity mice, 519

Fc receptor, rat:
preferential expression on red pulp macrophages, 329
FcR blocking activity:
in renal allograft recipients due to anti-class II MHC alloantibody, 379
Fimbrae:
immune response to, 8
Foot-and-mouth disease virus:
capsid protein epitopes represented by synthetic peptides, 1

Germinal centre formation:
regulation of, 487
Germinal centre T lymphocytes, 542

Hepatitis A:
clonal analysis of infiltrating T cells in liver, 14
Histamine:
regulation of IL-1 synthesis, 16
HRF20:
expression on peripheral blood mononuclear cells, 396

Idiotypes, shared, on DNA-binding antibodies, 348
IFN-γ, human:
up-regulates IL-2 receptors in activated T lymphocytes, 554
IL-1:
inhibition of production by placental protein 14, 277
IL-1, synthesis:
regulation by histamine, 162
IL-2:
control of autoimmune diabetes, 209
IL-2 activity:
in lymph fluid following *in vivo* antigen challenge, 45
IL-2 receptor:
decreased level of mRNA in tumour-bearing animals, 626

IL-2 synthesis:
role of intracellular calcium release and extracellular calcium influx, 468
IL-4:
induction of CD25 on B cells, 57
IL-6:
secretion by human monocytes, 52
IL-8:
in endothelial cells, 548
IgA:
inhibition by neutrophil chemotaxis, 257
IgA, polymeric:
response to typhoid vaccination, 190
IgE regulation:
role of $CD8^+$ and $CD4^+$ cells, 71
IgE responses, MHC-restricted:
to *Ascaris* antigen, 596
IgG4 antibodies to KLH in man, 355
Immunosuppression, viral:
inhibition of diabetes, 501

Junin virus:
induction of contrasuppressor cells, 629

L-leucyl-leucine-methyl ester:
toxicity for lymphocytes, 564
Leukotriene C_4:
release from human eosinophils, 435
Locomotion, chemotaxis and clustering:
of immune cells, 127
Ly-6:
characterization of monoclonal antibody to, 558
Lymphokine production:
synergism between AS101 and PMA, 110

Marmoset, immunoregulations in:
properties of lymphocytes, 616
Mast cell plasma membrane:
cleavage by chymase, 423
Mast cell proteinases:
interactions with serpins in blood, 139
Mast cell sensitization:
inhibition, 150
Membrane-bound hapten:
location, 323
MHC- and macrophage-associated antigen:
differential expression in fetal and postnatal small intestine, 409
MHC class II antigen:
expression in human vascular smooth muscle cells, 243
Mixed isotype $A_\beta E_\alpha$ class II molecules in transgenic mice, 385
Monocytic cells:
activation by antibodies CD11a/18 (LFA-1) complex, 574
Mononuclear cells, tumour associated:
in colonic carcinomas, 282
Mononuclear phagocytes, human:
differential effects of LPS, IFN-γ and TNFα on secretion of lysozyme, 402

MRL-lpr mice:
treatment with immunosuppressive FK506, 222
Mucosal immunity, bowel:
reactivity to virus infection, 27

Natural killer cells, human:
heterogeneity of, 166
Neutrophil adherence, to endothelium:
down-regulation by phorbol ester, 429
Non-lymphoid cells, rat thymic:
characterization, 416

Oncogene expression: in B lymphocytes, 490
Osteoblast-like cell line:
maintenance of mast cell viability and activity, 145

Phosphorylcholine antigen, from *T. spiralis*:
protection of mice from pneumococcus, 443
Popliteal lymphocytes:
ovine, deficient responsiveness, 134

Ricin:
sensitivity of rat $CD8^+$ and $CD4^+$ cells, 71

Salmonella strains:
aromatic or purine-dependent, immune responses to, 184
Skin grafts, H-Y disparate:
immunity to, without priming of CTL, 85
Suppressor cells:
inhibition of IL- 2 production, 454
Suppressor cells, antigen presentation to, by epidermal cells, 97
Suppressor-inducer capabilities:
associated with epitopes expressed on keratinocytes and Langerhans' cells, 622
Synovial cell, rheumatoid:
stimulation by complement, 237
Synthetic peptides:
induction of neutralizing antibodies to FMDV, 171
Systemic lupus erythrematosus, experimental:
genetic regulation, 228

T-cell receptor chains:
in vivo elimination in mice, 508
T cells, non-recirculating:
selective attrition, 476
TNF:
production by mastocytoma P815, 312
TNF and IL-1 and *Toxoplasma gondii*, 33
TNF in leishmaniasis, 570
TNFα:
potentiation of effect on anti-bacterial resistance by interleukin-1, 316
Thy-1:
molecular nature and cellular localization, 391
Thymocytes, murine:
growth in chemically defined medium, 303
Transgenic mice:
mixed isotype class II molecules, 385